

Supplementary Table 1. Percentages of tropical acute febrile illnesses patients by clinical characteristics ($n=336$).

	Leptospiros (%)	<i>P</i>	Dengue (%)	<i>P</i>	UAFI (%)	<i>P</i>	Bacteremia (%)	<i>P</i>	Melidosis (%)	<i>P</i>	Malaria (%)	<i>P</i>
AKI												
Yes	23.3	0.807	23.3	< .001	21.8	0.695	15	< .001	12	< .001	4.5	0.973
No	22.2		43.8		23.7		3.9		2		4.43	
AKI Stages												
AKI stage 1	22.4	0.924	35.5	< .001	21.1	0.96	11.8	< .001	4	< .001	5.3	0.698
AKI stage 2	27.3		6.1		24.2		9.1		27.3		6.1	
AKI stage 3	20.8		8.3		20.8		33.3		16.7		0	
Platelet count												
0 to 10	25	0.003	50	< .001	0	< .001	0	0.002	25	0.053	0	0.043
11 to 99	11.6		62.5		8.9		2.7		5.36		8.9	
100 to 150	35.6		42.4		15.3		3.4		0		3.4	
≥ 151	25.5		14.3		36		14.3		8.1		1.9	
RRT initiation												
Yes	17.2	0.469	10.3	0.003	13.8	0.221	27.6	< .001	31	< .001	0	0.223
No	23.1		38.1		23.8		6.5		3.6		4.89	
In-hospital mortality												
Yes	24.3	0.024	39	< .001	22.3	0.395	6.2	< .0001	3.3	< .001	4.9	0.206
No	6.5		3.23		29		29		32.3		0	

Notes: Significant at P -value < 0.05; *Chi*-square test statistics for categorical variables.